

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Rabbit anti Human Serum proteins, Clone: [Polyclonal], Monoclonal NMB-RAHU/TSP

|                            |  |
|----------------------------|--|
| Article Name               | Rabbit anti Human Serum proteins, Clone: [Polyclonal], Monoclonal  |
| Biozol Catalog Number      | NMB-RAHU/TSP   |
| Supplier Catalog Number    | RAHu/TSP   |
| Alternative Catalog Number | NMB-RAHU/TSP   |
| Manufacturer               | NordicMubio  |
| Host                       | Rabbit   |
| Category                   | Antikörper   |
| Product Description        | In immunoelectrophoresis against pooled serum and enriched serum proteins fractions precipitation can be observed of not less than 16 individual proteins components. In precipitating techniques as immunoelectrophoresis and radial immunodiffusion (Ouc...  |
| Clonality                  | Monoclonal   |
| Clone Designation          | [Polyclonal]   |
| Buffer                     | Delipidated, heat inactivated, lyophilized, stable whole antiserum No preservative added. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal rabbit serum. No foreign proteins added. Reconstitute the lyophilized |
| Source                     | Pooled whole human serum and partly purified serum fractions. Freund's complete adjuvant is used in the first step of the immunization procedure.  |

|                   |   |
|-------------------|---|
| Formula           | Delipidated, heat inactivated, lyophilized, stable whole antiserum No preservative added. Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal rabbit serum. No foreign proteins added.   |
| Application Notes | Precipitation assays. In immunoelectrophoresis use 2 $\mu$ l serum or equivalent against 120 $\mu$ l antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 $\mu$ l antiserum in 3 mm diameter center well and 2 $\mu$ l serum samples (neat |