

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## **Product Datasheet**

## Rabbit anti Mouse IgE (Fc specific), conjugated with TRITC, Clone: [Polyclonal], Monoclonal NMB-RAM/IGE(FC)/TRITC

| Article Name               | Rabbit anti Mouse IgE (Fc specific), conjugated with TRITC, Clone:<br>[Polyclonal], Monoclonal  |
|----------------------------|---|
| Biozol Catalog Number      | NMB-RAM/IGE(FC)/TRITC   |
| Supplier Catalog Number    | RAM/IgE(Fc)/TRITC   |
| Alternative Catalog Number | NMB-RAM/IGE(FC)/TRITC   |
| Manufacturer               | NordicMubio   |
| Host                       | Rabbit  |
| Category                   | Antikörper  |
| Conjugation                | TRITC   |
| Product Description        | The reactivity of the antiserum is directed to IgE as tested against<br>mouse sera and a panel of purified homogenous immunoglobulins.<br>The precipitation reaction obtained against serum containing<br>monoclonal IgE shows a reaction of full identity with the     |
| Clonality                  | Monoclonal  |
| Clone Designation          | [Polyclonal]  |
| Buffer                     | Fluorochrome-coupled purified hyperimmune rabbit IgG lyophilized<br>from a solution in phosphate buffered saline (PBS, pH 7.2). No<br>preservative added, as it may interfere with the antibody activity. It<br>is reconstituted by adding 1 ml sterile distilled water |

## BIOZOL

| Source            | Pools of highly purified homogenous IgE isolated from mouse serum.<br>Freunds complete adjuvant is used in the first step of the<br>immunization procedure.   |
|-------------------|---|
| Formula           | Fluorochrome-coupled purified hyperimmune rabbit IgG lyophilized<br>from a solution in phosphate buffered saline (PBS, pH 7.2). No<br>preservative added, as it may interfere with the antibody activity. |
| Application Notes | ELISA,Immunocytochemistry,Immunohistochemistry<br>(frozen),(In)direct immunofluorescence.   |