

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

**Telephone:** +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## **Product Datasheet**

## Rabbit anti Mouse IgG1 IgG2a IgG2b IgG3 (Fc specific), conjugated with Biotin, Clone: [Polyclonal], Monoclonal NMB-RAM/IGG(FC)/BIO

| Article Name               | Rabbit anti Mouse IgG1 IgG2a IgG2b IgG3 (Fc specific), conjugated with Biotin, Clone: [Polyclonal], Monoclonal   |
|----------------------------|--|
| Biozol Catalog Number      | NMB-RAM/IGG(FC)/BIO  |
| Supplier Catalog Number    | RAM/IgG(Fc)/Bio  |
| Alternative Catalog Number | NMB-RAM/IGG(FC)/BIO  |
| Manufacturer               | NordicMubio  |
| Host                       | Rabbit   |
| Category                   | Antikörper   |
| Conjugation                | Biotin   |
| Product Description        | The reactivity of the antiserum is directed to the Fc subunit of the IgG molecule. The antiserum contains antibodies to subclass-specific determinants as well as to determinants shared by two or more subclasses of IgG. In immunoelectrophoresis and do     |
| Clonality                  | Monoclonal   |
| Clone Designation          | [Polyclonal]   |
| Buffer                     | Biotin-coupled purified hyperimmune rabbit IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity. IgG protein concentration 10 mg/ml. Biotin/IgG protein molar rat |

| Source            | Purified normal IgG, including all known subclasses, isolated from pooled mouse serum. Freunds complete adjuvant is used in the first step of the immunization procedure.  |
|-------------------|--|
| Formula           | Biotin-coupled purified hyperimmune rabbit IgG lyophilized from a solution in phosphate buffered saline (PBS, pH 7.2). No preservative added, as it may interfere with the antibody activity. IgG protein concentration 10 mg/ml. Biotin/IgG protein molar rat |
| Application Notes | ELISA,Immunocytochemistry,Immunohistochemistry (paraffin),Dot blot,Immunoblotting.   |