

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Rabbit anti Mouse IgM (Fc specific), Clone: [Polyclonal], Monoclonal NMB-RAM/IGM(FC)

| | |
|----------------------------|---|
| Article Name | Rabbit anti Mouse IgM (Fc specific), Clone: [Polyclonal], Monoclonal |
| Biozol Catalog Number | NMB-RAM/IGM(FC) |
| Supplier Catalog Number | RAM/IgM(Fc) |
| Alternative Catalog Number | NMB-RAM/IGM(FC) |
| Manufacturer | NordicMubio |
| Host | Rabbit |
| Category | Antikörper |
| Product Description | The reactivity of the antiserum is restricted to the Fc part of the IgM molecule. In immunoelectrophoresis and radial immunodiffusion, using various antiserum concentrations against normal mouse serum a single precipitin line is obtained which shows ... |
| Clonality | Monoclonal |
| Clone Designation | [Polyclonal] |
| Buffer | Delipidated, heat inactivated, lyophilized, stable whole antiserum No preservative added Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal rabbit serum. No foreign proteins added. Reconstitute the lyophilized |
| Source | Highly purified homogenous IgM isolated from mouse serum. Freund's complete adjuvant is used in the first step of the immunization procedure. |

| | |
|-------------------|--|
| Formula | Delipidated, heat inactivated, lyophilized, stable whole antiserum No preservative added Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal rabbit serum. No foreign proteins added. |
| Application Notes | Precipitation assays. In immunoelectrophoresis use 2 μ l serum or equivalent against 120 μ l antiserum In double radial immunodiffusion use a rosette arrangement with 10 μ l antiserum in 3 mm diameter centre well and 2 μ l serum samples (neat and serially di |