

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Rabbit anti Swine albumin, Clone: [Polyclonal], Monoclonal NMB-RASW/ALB

|                            |  |
|----------------------------|--|
| Article Name               | Rabbit anti Swine albumin, Clone: [Polyclonal], Monoclonal   |
| Biozol Catalog Number      | NMB-RASW/ALB   |
| Supplier Catalog Number    | RASw/Alb   |
| Alternative Catalog Number | NMB-RASW/ALB   |
| Manufacturer               | NordicMubio  |
| Host                       | Rabbit   |
| Category                   | Antikörper   |
| Product Description        | The defined antibody specificity is directed to albumin as tested against swine sera. In immunoelectrophoresis and double radial immunodiffusion (Ouchterlony), using various antiserum concentrations against appropriate concentrations of the immunogen...  |
| Clonality                  | Monoclonal   |
| Clone Designation          | [Polyclonal]   |
| Buffer                     | Delipidated, heat inactivated, lyophilized, stable whole antiserum. No preservative added. Total protein and IgG concentrations in the antiserum are comparable to those in pooled rabbit serum. No foreign proteins added. Reconstitute the lyophilized artis |
| Source                     | Highly purified albumin isolated from swine serum. Freund's complete adjuvant is used in the first step of the immunization procedure.   |

|                   |  |
|-------------------|--|
| Formula           | Delipidated, heat inactivated, lyophilized, stable whole antiserum. No preservative added. Total protein and IgG concentrations in the antiserum are comparable to those in pooled rabbit serum. No foreign proteins added.  |
| Application Notes | Precipitation assays. In immunoelectrophoresis use 2 $\mu$ l serum, plasma or equivalent against 120 $\mu$ l antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 $\mu$ l antiserum in a 3 mm diameter centre well and 2 $\mu$ l serum samp |