

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Rabbit anti Swine IgM (Fc specific), Clone: [Polyclonal], Monoclonal NMB-RASW/IGM(FC)

| | |
|----------------------------|--|
| Article Name | Rabbit anti Swine IgM (Fc specific), Clone: [Polyclonal], Monoclonal |
| Biozol Catalog Number | NMB-RASW/IGM(FC) |
| Supplier Catalog Number | RASw/IgM(Fc) |
| Alternative Catalog Number | NMB-RASW/IGM(FC) |
| Manufacturer | NordicMubio |
| Host | Rabbit |
| Category | Antikörper |
| Product Description | The reactivity of the antiserum is restricted to the Fc part of the IgM molecule. In immunoelectrophoresis and radial immunodiffusion, using various antiserum concentrations against normal swine serum a single precipitin line is obtained which shows ... |
| Clonality | Monoclonal |
| Clone Designation | [Polyclonal] |
| Buffer | Delipidated, heat inactivated, lyophilized, stable whole antiserum. No preservative added Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal rabbit serum. No foreign proteins added. Reconstitute the lyophilized |
| Source | Highly purified normal IgM isolated from pooled swine serum. Freund's complete adjuvant is used in the first step of the immunization procedure. |

| | |
|-------------------|--|
| Formula | Delipidated, heat inactivated, lyophilized, stable whole antiserum. No preservative added Total protein and IgG concentrations in the antiserum are comparable to those of pooled normal rabbit serum. No foreign proteins added. |
| Application Notes | Precipitation assays. In immunoelectrophoresis use 2 μ l serum, plasma or equivalent against 120 μ l antiserum. In double radial immunodiffusion (Ouchterlony) use a rosette arrangement with 10 μ l antiserum in a 3 mm diameter centre well and 2 μ l serum samp |