

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

**Telephone:** +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## **Product Datasheet**

## GNAS Antibody, Rabbit, Polyclonal NSJ-F40026-0.4ML

| Article Name GNAS Antibody, Rabbit, Polyclonal  Biozol Catalog Number NSJ-F40026-0.4ML  Supplier Catalog Number F40026-0.4ML  Alternative Catalog Number NSJ-F40026-0.4ML  Manufacturer NSJ Bioreagents  |                 |
|--|-----------------|
| Supplier Catalog Number F40026-0.4ML  Alternative Catalog Number NSJ-F40026-0.4ML  |                 |
| Alternative Catalog Number NSJ-F40026-0.4ML  |                 |
|  |                 |
| Manufacturer NSJ Bioreagents   |                 |
|  |                 |
| Host Rabbit  |                 |
| Category Antikörper  |                 |
| Application ELISA, FACS, IF, IHC, WB   |                 |
| Species Reactivity Human   |                 |
| Immunogen A portion of amino acids 287-315 from the human protein as the immunogen for this GNAS antibody.   | n was used      |
| Product Description  Guanine nucleotide-binding proteins (G proteins) are involved modulators or transducers in various transmembrane sign systems. The Gs protein is involved in hormonal regulation adenylate cyclase: it activates the cyclase in response to | naling<br>on of |
| Clonality Polyclonal   |                 |
| Concentration In 1X PBS, pH 7.4, with 0.09% sodium azide   |                 |
| Isotype Rabbit Ig  |                 |
| UniProt Q5FWY2   |                 |
| Buffer In 1X PBS, pH 7.4, with 0.09% sodium azide  |                 |

| Purity             | Antigen affinity  |
|--------------------|---|
| Form               | In 1X PBS, pH 7.4, with 0.09% sodium azide  |
| Target             | GNAS  |
| Antibody Type      | Primary Antibody  |
| Application Dilute | Western blot: 1:1000,IHC (Paraffin): 1:10-1:50,Immunofluorescence: 1:10-1:50,Flow Cytometry: 1:10-1:50              |
| Application Notes  | Titration of the GNAS antibody may be required due to differences in protocols and secondary/substrate sensitivity. |