

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Golgi Antibody, Clone: [GLG1/970], CF488, Mouse, Monoclonal NSJ-V2543CF488-100T

| | |
|----------------------------|---|
| Article Name | Golgi Antibody, Clone: [GLG1/970], CF488, Mouse, Monoclonal |
| Biozol Catalog Number | NSJ-V2543CF488-100T |
| Supplier Catalog Number | V2543CF488-100T |
| Alternative Catalog Number | NSJ-V2543CF488-100T |
| Manufacturer | NSJ Bioreagents |
| Host | Mouse |
| Category | Antikörper |
| Application | FACS, IF |
| Species Reactivity | Human |
| Immunogen | The Golgi fraction from human liver cells was used as the immunogen for the Golgi Marker antibody. |
| Conjugation | CF488 |
| Product Description | This MAb recognizes a protein of 134kDa, which binds fibroblast growth factor and E-selectin (cell-adhesion lectin on endothelial cells mediating the binding of neutrophils). Fucosylation is essential for binding to E-selectin. It contains sialic aci... |
| Clonality | Monoclonal |
| Concentration | 500 ul at 0.1 mg/ml with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide |
| Clone Designation | [GLG1/970] |

| | |
|--------------------|--|
| Isotype | Mouse IgG1, kappa |
| UniProt | Q92896 |
| Buffer | 500 ul at 0.1 mg/ml with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide |
| Purity | Protein G affinity chromatography |
| Form | 500 ul at 0.1 mg/ml with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide |
| Target | Golgi |
| Antibody Type | Primary Antibody |
| Application Dilute | Flow cytometry: 5ul per test per one million cells in 0.1ml or 5ul per 100ul of whole blood,Immunofluorescence: 1-2ug/ml |
| Application Notes | Optimal dilution of the Golgi Marker antibody should be determined by the researcher. |