

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Fibrinogen alpha chain Antibody (FGA), Clone: [UC45], Mouse, Monoclonal NSJ-V7771SAF-100UG

| Article Name | Fibrinogen alpha chain Antibody (FGA), Clone: [UC45], Mouse, Monoclonal |
|----------------------------|---|
| Biozol Catalog Number | NSJ-V7771SAF-100UG |
| Supplier Catalog Number | V7771SAF-100UG |
| Alternative Catalog Number | NSJ-V7771SAF-100UG |
| Manufacturer | NSJ Bioreagents |
| Host | Mouse |
| Category | Antikörper |
| Application | ELISA, FACS, IF |
| Species Reactivity | Human |
| Immunogen | Human acute monoblastic leukemia cells were used as the immunogen for the Fibrinogen Alpha Chain antibody. |
| Product Description | The plasma glycoprotein Fibrinogen is synthesized in the liver and comprises three structurally different subunits: . Fibrinogen is important in platelet aggregation, the final step of the coagulation cascade (i.e. the formation of Fibrin) and determ |
| Clonality | Monoclonal |
| Concentration | 1 mg/ml in 1X PBS, BSA free, sodium azide free |
| Clone Designation | [UC45] |
| Isotype | Mouse IgM, kappa |

| Buffer | 1 mg/ml in 1X PBS, BSA free, sodium azide free |
|--------------------|---|
| Purity | PEG precipitation |
| Form | 1 mg/ml in 1X PBS, BSA free, sodium azide free |
| Target | Fibrinogen alpha chain |
| Antibody Type | Primary Antibody |
| Application Dilute | ELISA (order BSA-free format for coating),Flow cytometry: 1- 2ug/million cells in 0.1ml,Immunofluorescence: 1-2ug/ml |
| Application Notes | Optimal dilution of the antibody should be determined by the researcher. |