

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

HLA-DRB1 Antibody, Clone: [Bra30], Mouse, Monoclonal NSJ-V7844SAF-100UG

-DRB1 Antibody, Clone: [Bra30], Mouse, Monoclonal V7844SAF-100UG
44645 100110
44SAF-100UG
V7844SAF-100UG
Bioreagents
se
körper
S
nan
nan REH cells (a non-T, non-B leukemia cell line) were used as immunogen for the HLA-DRB1 antibody.
MAb reacts with a 28kDa chain of HLA-DRB1 antigen, a member HC class II molecules. It does not cross react with HLA-DP and -DQ. HLA-DR is a heterodimeric cell surface glycoprotein prised of a 36kDa alpha (heavy) chain and a 28kDa beta
oclonal
g/ml in 1X PBS, BSA free, sodium azide free
30]
se IgG2a, kappa
911

Buffer	1 mg/ml in 1X PBS, BSA free, sodium azide free
Purity	Protein G affinity chromatography
Form	1 mg/ml in 1X PBS, BSA free, sodium azide free
Target	HLA-DRB1
Antibody Type	Primary Antibody
Application Dilute	Flow cytometry: 1-2ug/million cells
Application Notes	Optimal dilution of the antibody should be determined by the researcher.