

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

MHC II I-Ad Antibody, Clone: [MK-D6], Mouse, Monoclonal NSJ-V8283-20UG

Article Name MHC II I-Ad Antibody, Clone: [MK-D6], Mouse, Monoclonal Biozol Catalog Number NSJ-V8283-20UG Supplier Catalog Number V8283-20UG Alternative Catalog Number NSJ-V8283-20UG Manufacturer NSJ Bioreagents Host Mouse Category Antikörper Application FACS, Functional Studies, IF Species Reactivity Mouse Immunogen Spleen cells from B10.D2 mice were used as the immunogen for the MHC II I-Ad antibody. Product Description Complex class II, I-Ad. It does not react with I-Ak, I-Ab, I-As, I-Af or I-Aa Clonality Monoclonal Concentration 0.2 mg/ml with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide Buffer 0.2 mg/ml with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide		
Supplier Catalog Number V8283-20UG Alternative Catalog Number NSJ-V8283-20UG Manufacturer NSJ Bioreagents Host Mouse Category Antikörper Application FACS, Functional Studies, IF Species Reactivity Mouse Immunogen Spleen cells from B10.D2 mice were used as the immunogen for the MHC II I-Ad antibody. Product Description The antibody is directed against Mouse major histocompatibility complex class II, I-Ad. It does not react with I-Ak, I-Ab, I-As, I-Af or I-Aa Clonality Monoclonal Concentration 0.2 mg/ml with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide Clone Designation [MK-D6] Isotype Mouse IgG2a, kappa	Article Name	MHC II I-Ad Antibody, Clone: [MK-D6], Mouse, Monoclonal
Alternative Catalog Number Manufacturer NSJ Bioreagents Host Mouse Category Antikörper Application FACS, Functional Studies, IF Species Reactivity Mouse Immunogen Spleen cells from B10.D2 mice were used as the immunogen for the MHC II I-Ad antibody. The antibody is directed against Mouse major histocompatibility complex class II, I-Ad. It does not react with I-Ak, I-Ab, I-As, I-Af or I-Aa Clonality Monoclonal Concentration 0.2 mg/ml with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide Clone Designation [MK-D6] Isotype Mouse IgG2a, kappa	Biozol Catalog Number	NSJ-V8283-20UG
Manufacturer NSJ Bioreagents Host Mouse Category Antikörper Application FACS, Functional Studies, IF Species Reactivity Mouse Immunogen Spleen cells from B10.D2 mice were used as the immunogen for the MHC II I-Ad antibody. Product Description The antibody is directed against Mouse major histocompatibility complex class II, I-Ad. It does not react with I-Ak, I-Ab, I-As, I-Af or I-Aa Clonality Monoclonal Concentration 0.2 mg/ml with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide Clone Designation [MK-D6] Isotype Mouse IgG2a, kappa	Supplier Catalog Number	V8283-20UG
Host Mouse Category Antikörper Application FACS, Functional Studies, IF Species Reactivity Mouse Immunogen Spleen cells from B10.D2 mice were used as the immunogen for the MHC II I-Ad antibody. The antibody is directed against Mouse major histocompatibility complex class II, I-Ad. It does not react with I-Ak, I-Ab, I-As, I-Af or I-Aa Clonality Monoclonal Concentration 0.2 mg/ml with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide Clone Designation [MK-D6] Isotype Mouse IgG2a, kappa	Alternative Catalog Number	NSJ-V8283-20UG
Category Antikörper Application FACS, Functional Studies, IF Species Reactivity Mouse Immunogen Spleen cells from B10.D2 mice were used as the immunogen for the MHC II I-Ad antibody. The antibody is directed against Mouse major histocompatibility complex class II, I-Ad. It does not react with I-Ak, I-Ab, I-As, I-Af or I-Aa Clonality Monoclonal Concentration 0.2 mg/ml with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide Clone Designation [MK-D6] Isotype Mouse IgG2a, kappa	Manufacturer	NSJ Bioreagents
Application FACS, Functional Studies, IF Species Reactivity Mouse Immunogen Spleen cells from B10.D2 mice were used as the immunogen for the MHC II I-Ad antibody. Product Description The antibody is directed against Mouse major histocompatibility complex class II, I-Ad. It does not react with I-Ak, I-Ab, I-As, I-Af or I-Aa Clonality Monoclonal Concentration 0.2 mg/ml with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide Clone Designation [MK-D6] Isotype Mouse IgG2a, kappa	Host	Mouse
Species Reactivity Mouse Spleen cells from B10.D2 mice were used as the immunogen for the MHC II I-Ad antibody. The antibody is directed against Mouse major histocompatibility complex class II, I-Ad. It does not react with I-Ak, I-Ab, I-As, I-Af or I-Aa Clonality Monoclonal Concentration 0.2 mg/ml with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide Clone Designation [MK-D6] Isotype Mouse IgG2a, kappa	Category	Antikörper
Immunogen Spleen cells from B10.D2 mice were used as the immunogen for the MHC II I-Ad antibody. The antibody is directed against Mouse major histocompatibility complex class II, I-Ad. It does not react with I-Ak, I-Ab, I-As, I-Af or I-Aa Clonality Monoclonal Concentration 0.2 mg/ml with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide Clone Designation [MK-D6] Isotype Mouse IgG2a, kappa	Application	FACS, Functional Studies, IF
MHC II I-Ad antibody. The antibody is directed against Mouse major histocompatibility complex class II, I-Ad. It does not react with I-Ak, I-Ab, I-As, I-Af or I-Aa Clonality Monoclonal Concentration 0.2 mg/ml with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide Clone Designation [MK-D6] Isotype Mouse IgG2a, kappa	Species Reactivity	Mouse
Product Description complex class II, I-Ad. It does not react with I-Ak, I-Ab, I-As, I-Af or I-Aa Clonality Monoclonal Concentration 0.2 mg/ml with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide Clone Designation [MK-D6] Isotype Mouse IgG2a, kappa	Immunogen	
Concentration 0.2 mg/ml with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide Clone Designation [MK-D6] Isotype Mouse IgG2a, kappa	Product Description	complex class II, I-Ad. It does not react with I-Ak, I-Ab, I-As, I-Af or I-
Clone Designation [MK-D6] Isotype Mouse IgG2a, kappa	Clonality	Monoclonal
Isotype Mouse IgG2a, kappa	Concentration	0.2 mg/ml with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide
	Clone Designation	[MK-D6]
Buffer 0.2 mg/ml with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	Isotype	Mouse IgG2a, kappa
	Buffer	0.2 mg/ml with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide

Purity	Protein G affinity chromatography
Form	0.2 mg/ml with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide
Target	MHC II I-Ad
Antibody Type	Primary Antibody
Application Dilute	Functional Studies (order BSA/azide-free format),Flow cytometry,Immunofluorescence
Application Notes	Optimal dilution of the antibody should be determined by the researcher.