

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Cig2 Antibody, Clone: [3A11/5], Mouse, Monoclonal NSJ-V8291-20UG

Cig2 Antibody, Clone: [3A11/5], Mouse, Monoclonal
NSJ-V8291-20UG
V8291-20UG
NSJ-V8291-20UG
NSJ Bioreagents
Mouse
Antikörper
ELISA, WB
Fungi
Bacterial S. pombe Cig2 purified protein was used as the immunogen for the Cig2 antibody.
Cig2 is a cyclin that functions as the catalytic subunit of cyclin- dependent kinases (Cdks) in the fission yeast Schizosaccharomyces pombe. It is a B-type, S phase cyclin that is required for both the G1/S and G2/M cell cycle transitions and is expre
Monoclonal
0.2 mg/ml with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide
[3A11/5]
Mouse IgG1, kappa
0.2 mg/ml with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide

Purity	Protein G affinity chromatography
Form	0.2 mg/ml with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide
Target	Cig2
Antibody Type	Primary Antibody
Application Dilute	ELISA (order BSA-free format for coating),Western blot: 1-2ug/ml
Application Notes	Optimal dilution of the antibody should be determined by the researcher.