

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

MECA-79 Antibody / Peripheral Node Addressin / PNAd, Clone: [MECA-79], Rat, Monoclonal NSJ-V8344SAF-100UG

| | |
|----------------------------|---|
| Article Name | MECA-79 Antibody / Peripheral Node Addressin / PNAd, Clone: [MECA-79], Rat, Monoclonal |
| Biozol Catalog Number | NSJ-V8344SAF-100UG |
| Supplier Catalog Number | V8344SAF-100UG |
| Alternative Catalog Number | NSJ-V8344SAF-100UG |
| Manufacturer | NSJ Bioreagents |
| Host | Rat |
| Category | Antikörper |
| Application | FACS, IF, IHC-P, WB |
| Species Reactivity | Human, Mouse, Rat |
| Immunogen | Mouse lymph node stromal cells were used as the immunogen for this MECA-79 antibody. |
| Product Description | MECA-79 recognizes a carbohydrate epitope shared with a group of sulfation decorated sialomucins, including sulfated ligands for CD62L (CD34, GlyCAM-1, Sgp200, and a subset of MAdCAM-1). This set antigens has been referred to as peripheral node addre... |
| Clonality | Monoclonal |
| Concentration | 1 mg/ml in 1X PBS, BSA free, sodium azide free |
| Clone Designation | [MECA-79] |
| Isotype | Rat IgM, kappa |

| | |
|--------------------|--|
| Buffer | 1 mg/ml in 1X PBS, BSA free, sodium azide free |
| Purity | Protein G affinity chromatography |
| Form | 1 mg/ml in 1X PBS, BSA free, sodium azide free |
| Target | MECA-79 |
| Antibody Type | Primary Antibody |
| Application Dilute | Flow cytometry: 1-2ug/million cells, Immunofluorescence: 1-2ug/ml, Western blot: 1-2ug/ml, Immunohistochemistry (FFPE): 1-2ug/ml |
| Application Notes | Optimal dilution of the antibody should be determined by the researcher. |