

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

**Telephone:** +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## **Product Datasheet**

CD4 Antibody, Clone: [GK1.5], Rat, Monoclonal

NSJ-V8352-100UG

Article Name CD4 Antibody, Clone: [GK1.5], Rat, Monoclonal  Biozol Catalog Number NSJ-V8352-100UG  Supplier Catalog Number V8352-100UG  Alternative Catalog Number NSJ-V8352-100UG  Manufacturer NSJ Bioreagents  Host Rat  Category Antikörper  Application ELISA, FACS, IF, NeA  Species Reactivity Mouse
Supplier Catalog Number V8352-100UG  Alternative Catalog Number NSJ-V8352-100UG  Manufacturer NSJ Bioreagents  Host Rat  Category Antikörper  Application ELISA, FACS, IF, NeA
Alternative Catalog Number NSJ-V8352-100UG  Manufacturer NSJ Bioreagents  Host Rat  Category Antikörper  Application ELISA, FACS, IF, NeA
Manufacturer  NSJ Bioreagents  Host  Rat  Category  Antikörper  Application  ELISA, FACS, IF, NeA
Host Rat  Category Antikörper  Application ELISA, FACS, IF, NeA
Category Antikörper  Application ELISA, FACS, IF, NeA
Application ELISA, FACS, IF, NeA
Species Reactivity Mouse
Immunogen Mouse CTL clone V4 was used as the immunogen for the CD4 antibody.
Product Description  CD4 is a member of the Ig superfamily, primarily expressed on most thymocytes, a subset of T cells, and weakly on macrophages and dendritic cells. It acts as a coreceptor with the TCR during T cell activation and thymic differentiation by binding MHC
Clonality Monoclonal
Concentration 0.2 mg/ml with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide
Clone Designation [GK1.5]
Isotype Rat IgG2b, kappa
UniProt P06332

Buffer	0.2 mg/ml with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide
Purity	Protein G affinity chromatography
Form	0.2 mg/ml with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide
Target	CD4
Antibody Type	Primary Antibody
Application Dilute	ELISA (order BSA-free format for coating & Damp; Azide), Neutralization Studies (order BSA/azide-free format), Flow cytometry: 1-2ug/million cells in 0.1ml, Immunofluorescence: 1-2ug/ml
Application Notes	Optimal dilution of the antibody should be determined by the researcher.