

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### CD20 Antibody / MS4A1, Clone: [109-3C2], Mouse, Monoclonal NSJ-V8375-100UG

|                            |   |
|----------------------------|---|
| Article Name               | CD20 Antibody / MS4A1, Clone: [109-3C2], Mouse, Monoclonal  |
| Biozol Catalog Number      | NSJ-V8375-100UG   |
| Supplier Catalog Number    | V8375-100UG   |
| Alternative Catalog Number | NSJ-V8375-100UG   |
| Manufacturer               | NSJ Bioreagents   |
| Host                       | Mouse   |
| Category                   | Antikörper  |
| Application                | ELISA, Functional Studies   |
| Species Reactivity         | Human   |
| Immunogen                  | Stimulated human leukocytes were used as the immunogen for the CD20 antibody.   |
| Product Description        | Recognizes a protein of 30-33kDa, which is identified as CD20 (Workshop V, Code CD20.12. Workshop IV, Code B17). It recognizes an extracellular domain of CD20. It is a non-Ig differentiation antigen of B-cells and its expression is restricted to norm... |
| Clonality                  | Monoclonal  |
| Concentration              | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide   |
| Clone Designation          | [109-3C2]   |
| Isotype                    | Mouse IgG3, kappa   |

|                    |  |
|--------------------|--|
| UniProt            | <a href="#">P11836</a>   |
| Buffer             | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide          |
| Purity             | Protein G affinity chromatography  |
| Form               | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide          |
| Target             | CD20   |
| Antibody Type      | Primary Antibody   |
| Application Dilute | Functional Studies: order Ab without BSA,ELISA: order Ab without BSA for coating |
| Application Notes  | Optimal dilution of the CD20 antibody should be determined by the researcher.    |