

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | Fax: +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

CD63 Antibody / LAMP-3, Clone: [LAMP3/4949], Mouse, Monoclonal NSJ-V8875SAF-100UG

CD63 Antibody / LAMP-3, Clone: [LAMP3/4949], Mouse, Monoclonal
NSJ-V8875SAF-100UG
V8875SAF-100UG
NSJ-V8875SAF-100UG
NSJ Bioreagents
Mouse
Antikörper
ELISA, FACS, IF, IHC-P
Human
Full-length human CD63 protein was used as the immunogen for the CD63 antibody.
CD63 is expressed on activated platelets, monocytes and macrophages, and is weakly expressed on granulocytes, T cell and B cells. It is located on the basophilic granule membranes and on the plasma membranes of lymphocytes and granulocytes. CD63 is a
Monoclonal
1 mg/ml in 1X PBS, BSA free, sodium azide free
[LAMP3/4949]
Mouse IgG1, kappa
P08962

Buffer	1 mg/ml in 1X PBS, BSA free, sodium azide free
Purity	Protein A/G affinity
Form	1 mg/ml in 1X PBS, BSA free, sodium azide free
Target	CD63
Antibody Type	Primary Antibody
Application Dilute	ELISA: 1-5ug/ml (order BSA-free format for coating),Flow cytometry: 1-2ug/million cells,Immunofluorescence: 1-2ug/ml,Immunohistochemistry (FFPE): 1-2ug/ml
Application Notes	Optimal dilution of the CD63 antibody should be determined by the researcher.