

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Recombinant p21WAF1 (Tumor Suppressor Protein) Antibody, Clone: [CIP1/2275R], Rabbit, Monoclonal NBT-1026-RBM7-P0

| | |
|----------------------------|---|
| Article Name | Recombinant p21WAF1 (Tumor Suppressor Protein) Antibody, Clone: [CIP1/2275R], Rabbit, Monoclonal |
| Biozol Catalog Number | NBT-1026-RBM7-P0 |
| Supplier Catalog Number | 1026-RBM7-P0 |
| Alternative Catalog Number | NBT-1026-RBM7-P0-20,NBT-1026-RBM7-P0-100 |
| Manufacturer | NeoBiotechnologies |
| Host | Rabbit |
| Category | Antikörper |
| Application | IHC, WB |
| Species Reactivity | Human |
| Immunogen | Recombinant full-length human CDKN1A protein |
| Product Description | This MAb recognizes a 21kDa protein, identified as the p21WAF1 tumor suppressor protein. This MAb is highly specific to p21 and shows no cross-reaction with other closely related mitotic inhibitors. p21WAF1 is a specific inhibitor of cdks and a tumor... |
| Clonality | Monoclonal |
| Clone Designation | [CIP1/2275R] |
| Molecular Weight | 21kDa |
| NCBI | 1026 |

| | |
|-------------------|---|
| UniProt | P38936 |
| Form | 200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Recombinant Monoclonal Antibody |
| Application Notes | Western Blot (1-2ug/ml), Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 min |