

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Recombinant CDX2 / Caudal Type Homeobox 2 (GI Epithelial Marker) Antibody, Clone: [CDX2/2951R], Rabbit, Monoclonal NBT-1045-RBM5-P1ABX

Article Name	Recombinant CDX2 / Caudal Type Homeobox 2 (GI Epithelial Marker) Antibody, Clone: [CDX2/2951R], Rabbit, Monoclonal
Biozol Catalog Number	NBT-1045-RBM5-P1ABX
Supplier Catalog Number	1045-RBM5-P1ABX
Alternative Catalog Number	NBT-1045-RBM5-P1ABX-100
Manufacturer	NeoBiotechnologies
Host	Rabbit
Category	Antikörper
Application	ELISA, IHC
Species Reactivity	Human
Immunogen	Recombinant fragment (around aa150-249) of human CDX2 protein (exact sequence is proprietary)
Product Description	The intestine-specific transcription factors CDX1 and CDX2 are important for directing intestinal development, differentiation, proliferation and maintenance of the intestinal phenotype. CDX2 protein expression has been seen in GI carcinomas. Anti-CD
Clonality	Monoclonal
Clone Designation	[CDX2/2951R]
Molecular Weight	40kDa
NCBI	1045

UniProt	Q99626
Form	200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double are applied as a solution of the protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double are a solution of the protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double are a solution of the protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double are a solution of the protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double are a solution of the protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double are a solution of the protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double are a solution of the protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double are a solution of the protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double are a solution of the protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double are a solution of the protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double are a solution of the protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double are a solution of the protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double are a solution of the protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double are a solution of the protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double are a solution of the protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double are a solution of the protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double are a solution of the protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double are a solution of the prepared in 10mM PBS with 0.05% BSA & Double are a solution of the prepared in 10mM PBS with 0.05% BSA & Double are a solution of the prepared in 10mM PBS with 0.05% BSA & Double are a solution of the prepared in 10mM PBS with 0.05% BSA & Double are a solution of the prepared in 10mM PBS with 0.05% BSA & Double are a solution of the prepared in 10mM PBS with 0.05% BSA & Double are a solution of the prepared in 10mM PBS with 0.05% BSA & Double are a solution of the 10mM PBS with 0.05% BSA & Double are a solution of the 10mM PBS with 0.
Antibody Type	Recombinant Monoclonal Antibody
Application Notes	ELISA (Use Ab at 2-4ug/ml for coating) (Order Ab without BSA), Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 min at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C