

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

**Telephone:** +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## **Product Datasheet**

## Calponin-1 (Smooth Muscle Marker) Antibody, IgG1, Clone: [CNN1/832 + CALP], Mouse, Monoclonal NBT-1264-MSM3-P0

| Article Name               | Calponin-1 (Smooth Muscle Marker) Antibody, IgG1, Clone: [CNN1/832 + CALP], Mouse, Monoclonal  |
|----------------------------|--|
| Biozol Catalog Number      | NBT-1264-MSM3-P0   |
| Supplier Catalog Number    | 1264-MSM3-P0   |
| Alternative Catalog Number | NBT-1264-MSM3-P0-20,NBT-1264-MSM3-P0-100   |
| Manufacturer               | NeoBiotechnologies   |
| Host                       | Mouse  |
| Category                   | Antikörper   |
| Application                | FC, IF, IHC  |
| Species Reactivity         | Human, Rat   |
| Immunogen                  | Recombinant full-length human CNN1 protein (CNN1/832), Crude human uterus extract (CALP)   |
| Product Description        | Multiple isoelectric variants of calponin have been identified, however only two molecular weight isoforms exist, a 34kDa form and a 29kDa form. Expression of the 29kDa form, I-calponin, is primarily restricted to muscle of the urogenital tract, wher |
| Clonality                  | Monoclonal   |
| Clone Designation          | [CNN1/832 + CALP]  |
| Molecular Weight           | 34kDa  |
| Isotype                    | IgG1   |

| NCBI              | 1264  |
|-------------------|---|
| UniProt           | P51911  |
| Form              | 200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double are a side. Also available WITHOUT BSA & Double at 1.0mg/ml.   |
| Antibody Type     | Monoclonal Antibody   |
| Application Notes | Flow Cytometry (1-2ug/million cells), Immunofluorescence (1-2ug/ml), Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 1mM EDTA, pH 7.5-8.5, for 45 min at 95&degC |