

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Recombinant CD21 (Mature B-Cell & Follicular Dendritic Cell Marker) Antibody, Clone: [CR2/3990R], Rabbit, Monoclonal NBT-1380-RBM12-P0

| | |
|----------------------------|---|
| Article Name | Recombinant CD21 (Mature B-Cell & Follicular Dendritic Cell Marker) Antibody, Clone: [CR2/3990R], Rabbit, Monoclonal |
| Biozol Catalog Number | NBT-1380-RBM12-P0 |
| Supplier Catalog Number | 1380-RBM12-P0 |
| Alternative Catalog Number | NBT-1380-RBM12-P0-20,NBT-1380-RBM12-P0-100 |
| Manufacturer | NeoBiotechnologies |
| Host | Rabbit |
| Category | Antikörper |
| Application | IHC |
| Species Reactivity | Human |
| Immunogen | Synthetic peptide corresponding to CD21 residues within aa933-C terminal of CD21 |
| Product Description | CD21 also known as complement C3d receptor, Epstein-Barr virus receptor, and complement receptor type 2 (CR2), is a protein that in humans is encoded by the CR2 gene. This protein is involved in the complement system. It binds to iC3b (inactive deriv... |
| Clonality | Monoclonal |
| Clone Designation | [CR2/3990R] |
| Molecular Weight | 140kDa |
| NCBI | 1380 |

| | |
|-------------------|---|
| UniProt | P20023 |
| Form | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Recombinant Monoclonal Antibody |
| Application Notes | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95 °C followed by cooling at RT for 20 minutes),Optimal diluti |