

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | Fax: +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Cystatin A Antibody, IgG2b, Clone: [CSTA/3553], Mouse, Monoclonal NBT-1475-MSM3-P1ABX

Cystatin A Antibody, IgG2b, Clone: [CSTA/3553], Mouse, Monoclonal
NBT-1475-MSM3-P1ABX
1475-MSM3-P1ABX
NBT-1475-MSM3-P1ABX-100
NeoBiotechnologies
Mouse
Antikörper
IHC, WB
Human
Recombinant fragment of human CSTA protein (around aa 1-98) (exact sequence is proprietary)
The cystatin superfamily encompasses proteins that contain multiple cystatin-like sequences. Some of the members are active cysteine protease inhibitors, while others have lost or perhaps never acquired this inhibitory activity. There are three inhib
Monoclonal
[CSTA/3553]
11kDa
lgG2b
1475

UniProt	P01040
Form	200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double are a side. Also available WITHOUT BSA & Double at 1.0mg/ml.
Antibody Type	Monoclonal Antibody
Application Notes	Western Blot (2-4ug/ml), Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20