

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Desmoglein-3 (Squamous Cell Marker) Antibody, IgG1, Clone: [DSG3/2838], Mouse, Monoclonal NBT-1830-MSM8-P0

| | |
|----------------------------|---|
| Article Name | Desmoglein-3 (Squamous Cell Marker) Antibody, IgG1, Clone: [DSG3/2838], Mouse, Monoclonal |
| Biozol Catalog Number | NBT-1830-MSM8-P0 |
| Supplier Catalog Number | 1830-MSM8-P0 |
| Alternative Catalog Number | NBT-1830-MSM8-P0-20,NBT-1830-MSM8-P0-100 |
| Manufacturer | NeoBiotechnologies |
| Host | Mouse |
| Category | Antikörper |
| Application | IHC |
| Species Reactivity | Human |
| Immunogen | Recombinant fragment (around aa 379-491) human DSG3 protein (exact sequence is proprietary) |
| Product Description | Recognizes a protein of 130kDa, identified as Desmoglein-3 (DSG3). This MAb is highly specific to Desmoglein-3 and does not cross-react with other members of the Desmoglein-family. DSG3 is a calcium-binding transmembrane glycoprotein component of des... |
| Clonality | Monoclonal |
| Clone Designation | [DSG3/2838] |
| Molecular Weight | 130kDa |
| Isotype | IgG1 |

| | |
|-------------------|---|
| NCBI | 1830 |
| UniProt | P32926 |
| Form | 200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Monoclonal Antibody |
| Application Notes | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes), Optimal dilution f |