

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

**Telephone:** +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## **Product Datasheet**

## Erythropoietin (EPO) (Marker of Placentation Disorders) Antibody, IgG1, Clone: [EPO/1367], Mouse, Monoclonal NBT-2056-MSM1-P1ABX

Article Name	Erythropoietin (EPO) (Marker of Placentation Disorders) Antibody, IgG1, Clone: [EPO/1367], Mouse, Monoclonal
Biozol Catalog Number	NBT-2056-MSM1-P1ABX
Supplier Catalog Number	2056-MSM1-P1ABX
Alternative Catalog Number	NBT-2056-MSM1-P1ABX-100
Manufacturer	NeoBiotechnologies
Host	Mouse
Category	Antikörper
Application	ELISA, IHC
Species Reactivity	Human
Immunogen	Recombinant fragment (around aa 28-162) of human EPO protein (exact sequence is proprietary)
Product Description	Recognizes a protein of about 37kDa, which is identified as Erythropoietin (EPO). Erythropoietin is a secreted, glycosylated cytokine hormone composed of four alpha helical bundles. It is the primary factor responsible for regulating erythropoiesis d
Clonality	Monoclonal
Clone Designation	[EPO/1367]
Molecular Weight	37kDa
Isotype	lgG1

NCBI	2056
UniProt	P01588
Form	200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double are a side. Also available WITHOUT BSA & Double at 1.0mg/ml.
Antibody Type	Monoclonal Antibody
Application Notes	ELISA (Use Ab at 2-4ug/ml for coating) (Order Ab without BSA), Immunohistochemistry (Formalin-fixed) (2-4ug/ml for 30 min at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C fo