

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Recombinant ERG (Vascular Endothelial & Prostate Marker) Antibody, IgG1, Clone: [rERG/6843], Mouse, Monoclonal NBT-2078-MSM3-P0

| Article Name | Recombinant ERG (Vascular Endothelial & Prostate Marker) Antibody, IgG1, Clone: [rERG/6843], Mouse, Monoclonal |
|----------------------------|--|
| Biozol Catalog Number | NBT-2078-MSM3-P0 |
| Supplier Catalog Number | 2078-MSM3-P0 |
| Alternative Catalog Number | NBT-2078-MSM3-P0-20,NBT-2078-MSM3-P0-100 |
| Manufacturer | NeoBiotechnologies |
| Host | Mouse |
| Category | Antikörper |
| Application | IHC |
| Species Reactivity | Human |
| Immunogen | Recombinant fragment (around aa450 to 479) (C-terminus) of human ERG (exact sequence is proprietary) |
| Product Description | ERG (ETS-related gene) is a proto-oncogene, a member of the ETS family of transcription factors. The ERG gene encodes for a nuclear protein, also called ERG, which is involved in hematopoietic and endothelial development. ERG remains constitually exp |
| Clonality | Monoclonal |
| Clone Designation | [rERG/6843] |
| Molecular Weight | 55kDa |
| Isotype | lgG1 |

| NCBI | 2078 |
|-------------------|---|
| UniProt | P11308 |
| Form | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & amp; 0.05% azide. Also available WITHOUT BSA & amp; azide at 1.0mg/ml. |
| Antibody Type | Recombinant Monoclonal Antibody |
| Application Notes | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95 °C followed by cooling at RT for 20 minutes)Optimal dilution |