

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

FOLH1 / PSMA (Prostate Epithelial Marker) Antibody, IgG2b, Clone: [FOLH1/2354], Mouse, Monoclonal NBT-2346-MSM2-P0

| Article Name | FOLH1 / PSMA (Prostate Epithelial Marker) Antibody, IgG2b, Clone: [FOLH1/2354], Mouse, Monoclonal |
|----------------------------|--|
| Biozol Catalog Number | NBT-2346-MSM2-P0 |
| Supplier Catalog Number | 2346-MSM2-P0 |
| Alternative Catalog Number | NBT-2346-MSM2-P0-20,NBT-2346-MSM2-P0-100 |
| Manufacturer | NeoBiotechnologies |
| Host | Mouse |
| Category | Antikörper |
| Application | IHC, WB |
| Species Reactivity | Canine, Equine, Feline, Human |
| Immunogen | Recombinant human FOLH1 protein fragment (around aa 232-433) (exact sequence is proprietary) |
| Product Description | Folate hydrolase 1 (FOLH1), also known as Prostate-specific membrane antigen (PSMA), is a type II transmembrane glycoprotein belonging to the M28 peptidase family. FOLH1 has two enzymatic activities, one as a prostate-specific integral membrane folat |
| Clonality | Monoclonal |
| Clone Designation | [FOLH1/2354] |
| Molecular Weight | 100kDa |
| Isotype | lgG2b |

| NCBI | 2346 |
|-------------------|--|
| UniProt | Q04609 |
| Form | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double are applied to the substitution of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double are applied to the substitution of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double are applied to the substitution of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double are applied to the substitution of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double are applied to the substitution of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double are applied to the substitution of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double are applied to the substitution of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double are applied to the substitution of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double are applied to the substitution of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double are applied to the substitution of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double are applied to the substitution of Ab purified from Bioreactor Concentrate by Protein A/G. P |
| Antibody Type | Monoclonal Antibody |
| Application Notes | Western Blotting (1-2ug/ml),,Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT fo |