

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Recombinant Anti Mullerian Hormone (AMH) / Mullerian Inhibiting Substance (MIS) Antibody, Clone: [AMH/6713R], Rabbit, Monoclonal NBT-268-RBM3-P0

|                            |   |
|----------------------------|---|
| Article Name               | Recombinant Anti Mullerian Hormone (AMH) / Mullerian Inhibiting Substance (MIS) Antibody, Clone: [AMH/6713R], Rabbit, Monoclonal  |
| Biozol Catalog Number      | NBT-268-RBM3-P0   |
| Supplier Catalog Number    | 268-RBM3-P0   |
| Alternative Catalog Number | NBT-268-RBM3-P0-20,NBT-268-RBM3-P0-100  |
| Manufacturer               | NeoBiotechnologies  |
| Host                       | Rabbit  |
| Category                   | Antikörper  |
| Application                | IHC   |
| Species Reactivity         | Human   |
| Immunogen                  | Synthetic peptide from the C-terminus of human AMH (aa 460-560) (exact sequence is proprietary)   |
| Product Description        | The transforming growth factor (TGF) superfamily is composed of numerous growth and differentiation factors, including TGF1-3, Mullerian inhibiting substance (MIS), growth/differentiation factor (GDF) 1-9, bone morphogenic protein (BMP) 2-8, glial ce... |
| Clonality                  | Monoclonal  |
| Clone Designation          | [AMH/6713R]   |
| Molecular Weight           | 70kDa (reduced), 140kDa (nonreduced)  |
| NCBI                       | <a href="#">268</a>   |

|                   |   |
|-------------------|---|
| UniProt           | <a href="#">P03971</a>  |
| Form              | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.  |
| Antibody Type     | Recombinant Monoclonal Antibody   |
| Application Notes | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 min at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95 °C followed by cooling at RT for 20 minutes),Optimal dilution f |