

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

**Telephone:** +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## **Product Datasheet**

## HSP90AB1 (Heat Shock Protein 90) Antibody, IgG1, Clone: [HSP90AB1/3953], Mouse, Monoclonal NBT-3326-MSM3-P0

| Article Name               | HSP90AB1 (Heat Shock Protein 90) Antibody, IgG1, Clone: [HSP90AB1/3953], Mouse, Monoclonal   |
|----------------------------|--|
| Biozol Catalog Number      | NBT-3326-MSM3-P0   |
| Supplier Catalog Number    | 3326-MSM3-P0   |
| Alternative Catalog Number | NBT-3326-MSM3-P0-20,NBT-3326-MSM3-P0-100   |
| Manufacturer               | NeoBiotechnologies   |
| Host                       | Mouse  |
| Category                   | Antikörper   |
| Application                | IHC, WB  |
| Species Reactivity         | Human  |
| Immunogen                  | Recombinant human HSP90AB1 protein fragment (around aa581-704) (exact sequence is proprietary)   |
| Product Description        | This gene encodes a member of the heat shock protein 90 family, these proteins are involved in signal transduction, protein folding and degradation and morphological evolution. This gene encodes the constitutive form of the cytosolic 90 kDa heat-shoc |
| Clonality                  | Monoclonal   |
| Clone Designation          | [HSP90AB1/3953]  |
| Molecular Weight           | ~90kDa   |
| Isotype                    | lgG1   |

| NCBI              | 3326  |
|-------------------|---|
| UniProt           | P08238  |
| Form              | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with0.05% BSA & D.05% azide. Also available WITHOUT BSA at 1.0mg/ml.   |
| Antibody Type     | Monoclonal Antibody   |
| Application Notes | Western Blot (1-2ug/ml), ,Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95 &degC followed by cooling at RT for |