

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Recombinant Cytokeratin 7 (Glandular and Transitional Epithelial Marker) Antibody, Clone: [KRT7/8121R], Rabbit, Monoclonal NBT-3855-RBM17-P1ABX

| | |
|----------------------------|---|
| Article Name | Recombinant Cytokeratin 7 (Glandular and Transitional Epithelial Marker) Antibody, Clone: [KRT7/8121R], Rabbit, Monoclonal |
| Biozol Catalog Number | NBT-3855-RBM17-P1ABX |
| Supplier Catalog Number | 3855-RBM17-P1ABX |
| Alternative Catalog Number | NBT-3855-RBM17-P1ABX-100 |
| Manufacturer | NeoBiotechnologies |
| Host | Rabbit |
| Category | Antikörper |
| Application | IHC |
| Species Reactivity | Human |
| Immunogen | Recombinant fragment (around aa1-200) of human KRT7 protein (exact sequence is proprietary) |
| Product Description | Cytokeratin 7 in humans is a protein encoded by the KRT7 gene. It is always as a keratin pair with Cytokeratin 19. Cytokeratin 7 is expressed on most ductal and glandular epithelium including lung, breast, bladder and female genital tract, but not in... |
| Clonality | Monoclonal |
| Clone Designation | [KRT7/8121R] |
| Molecular Weight | 55kDa |
| NCBI | 3855 |

| | |
|-------------------|--|
| UniProt | P08729 |
| Form | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Recombinant Monoclonal Antibody |
| Application Notes | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes),Optimal dilutio |