

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

LH-beta (Luteinizing Hormone-beta) Antibody, IgG1, Clone: [LHb/1214], Mouse, Monoclonal NBT-3972-MSM1-P1ABX

| | |
|----------------------------|---|
| Article Name | LH-beta (Luteinizing Hormone-beta) Antibody, IgG1, Clone: [LHb/1214], Mouse, Monoclonal |
| Biozol Catalog Number | NBT-3972-MSM1-P1ABX |
| Supplier Catalog Number | 3972-MSM1-P1ABX |
| Alternative Catalog Number | NBT-3972-MSM1-P1ABX-100 |
| Manufacturer | NeoBiotechnologies |
| Host | Mouse |
| Category | Antikörper |
| Application | IHC |
| Species Reactivity | Human |
| Immunogen | Recombinant beta sub-unit of human LH |
| Product Description | Luteinizing hormone (LH) is a glycoprotein. Each monomeric unit is a sugar-like protein molecule, two of these make the full, functional protein. Its structure is similar to the other glycoproteins, follicle-stimulating hormone (FSH), thyroid-stimula... |
| Clonality | Monoclonal |
| Clone Designation | [LHb/1214] |
| Molecular Weight | 22kDa |
| Isotype | IgG1 |

| | |
|-------------------|---|
| NCBI | 3972 |
| UniProt | P01229 |
| Form | 200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Monoclonal Antibody |
| Application Notes | Immunohistochemistry (Formalin-fixed) (1-2ug/ml, 30 min RT) (Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes), Optimal dilution for a spec |