

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

**Telephone:** +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## **Product Datasheet**

## EpCAM / CD326 (Epithelial Marker) Antibody, IgG1, Clone: [Cocktail PAN-EpCAM], Mouse, Monoclonal NBT-4072-MSM9-P0

Article Name	EpCAM / CD326 (Epithelial Marker) Antibody, IgG1, Clone: [Cocktail PAN-EpCAM], Mouse, Monoclonal
Biozol Catalog Number	NBT-4072-MSM9-P0
Supplier Catalog Number	4072-MSM9-P0
Alternative Catalog Number	NBT-4072-MSM9-P0-20,NBT-4072-MSM9-P0-100
Manufacturer	NeoBiotechnologies
Host	Mouse
Category	Antikörper
Application	FC, IF, IHC, WB
Species Reactivity	Human
Immunogen	Recombinant human EpCAM protein (full-length and fragments)
Product Description	It is a cocktail of four highly specific monoclonal antibodies that recognize extracellular as well as intracellular domains of the epithelial cellular adhesion molecule (EpCAM). It is a 40-43kDa transmembrane epithelial glycoprotein, identified as e
Clonality	Monoclonal
Clone Designation	[Cocktail PAN-EpCAM]
Molecular Weight	40-43kDa
Isotype	lgG1

NCBI	4072
UniProt	P16422
Form	200ug/ml of Abs purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double are a side. Also available WITHOUT BSA & Double at 1.0mg/ml.
Antibody Type	Monoclonal Antibody
Application Notes	Flow Cytometry (1-2ug/million cells), Immunofluorescence (1-2ug/ml), Western Blot (1-2ug/ml), ,Immunohistochemistry (Formalinfixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1m