

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Recombinant SMAD4 / DPC4 (Pancreatic Adenocarcinoma Marker / Tumor Suppressor) Antibody, Clone: [SMAD/7905R], Rabbit, Monoclonal NBT-4089-RBM25-P1ABX

Article Name	Recombinant SMAD4 / DPC4 (Pancreatic Adenocarcinoma Marker / Tumor Suppressor) Antibody, Clone: [SMAD/7905R], Rabbit, Monoclonal
Biozol Catalog Number	NBT-4089-RBM25-P1ABX
Supplier Catalog Number	4089-RBM25-P1ABX
Alternative Catalog Number	NBT-4089-RBM25-P1ABX-100
Manufacturer	NeoBiotechnologies
Host	Rabbit
Category	Antikörper
Application	IHC
Species Reactivity	Human
Immunogen	Recombinant fragment (around aa400-552) of human SMAD4 protein (exact sequence is proprietary)
Product Description	Signaling from the ligand-activated membrane receptor serine/threonine kinases to nuclear targets is mediated by a set of evolutionarily conserved proteins known as DPC4. Upon ligand binding, the receptors of the TGF-i family phosphorylate SMAD prot
Clonality	Monoclonal
Clone Designation	[SMAD/7905R]
Molecular Weight	61kDa

NCBI	4089
UniProt	Q13485
Form	200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prote
Antibody Type	Recombinant Monoclonal Antibody
Application Notes	Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes),Optimal dilutio