

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## **Product Datasheet**

## MCAM (Melanoma Cell Adhesion Molecule) / MUC18 / CD146 Antibody, IgG1, Clone: [C146/634], Mouse, Monoclonal NBT-4162-MSM3-P1ABX

| Article Name               | MCAM (Melanoma Cell Adhesion Molecule) / MUC18 / CD146<br>Antibody, IgG1, Clone: [C146/634], Mouse, Monoclonal   |
|----------------------------|--|
| Biozol Catalog Number      | NBT-4162-MSM3-P1ABX  |
| Supplier Catalog Number    | 4162-MSM3-P1ABX  |
| Alternative Catalog Number | NBT-4162-MSM3-P1ABX-100  |
| Manufacturer               | NeoBiotechnologies   |
| Host                       | Mouse  |
| Category                   | Antikörper   |
| Application                | FC, IF, IHC  |
| Species Reactivity         | Human  |
| Immunogen                  | Recombinant human CD146 protein  |
| Product Description        | The human CD146 gene maps to chromosome 11q23 and encodes a trans-membrane glycoprotein, also designated MCAM, MUC 18, that belongs to the immunoglobulin superfamily and functions as a Ca2+-independent cell adhesion molecule. CD146 expression is rest |
| Clonality                  | Monoclonal   |
| Clone Designation          | [C146/634]   |
| Molecular Weight           | 130kDa   |
| Isotype                    | lgG1   |

| NCBI              | 4162  |
|-------------------|---|
| UniProt           | P43121  |
| Form              | 200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G.<br>Prepared in 10mM PBS with 0.05% BSA & amp; 0.05% azide. Also<br>available WITHOUT BSA & amp; azide at 1.0mg/ml.  |
| Antibody Type     | Monoclonal Antibody   |
| Application Notes | Flow Cytometry (1-2ug/million cells), Immunofluorescence (1-<br>2ug/ml), Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30<br>minutes at RT)(Staining of formalin-fixed tissues requires heating<br>tissue sections in 1mM EDTA, pH 7.5-8.5, for 45 min at 95&degC |