

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Recombinant c-Myc Oncoprotein Antibody, Clone: [MYC/7150R], Rabbit, Monoclonal NBT-4609-RBM10-P0

| | |
|----------------------------|---|
| Article Name | Recombinant c-Myc Oncoprotein Antibody, Clone: [MYC/7150R], Rabbit, Monoclonal |
| Biozol Catalog Number | NBT-4609-RBM10-P0 |
| Supplier Catalog Number | 4609-RBM10-P0 |
| Alternative Catalog Number | NBT-4609-RBM10-P0-20,NBT-4609-RBM10-P0-100 |
| Manufacturer | NeoBiotechnologies |
| Host | Rabbit |
| Category | Antikörper |
| Application | IHC |
| Species Reactivity | Human |
| Immunogen | Recombinant fragment of human MYC protein (exact sequence is proprietary) |
| Product Description | It recognizes a transcription factor of 64-67kDa, identified as c-myc. Its epitope spans between aa 410-419 (EQKLISEEDL) which is a specific portion of an alpha helical region of human c-myc protein. This MAb shows no cross-reaction with v-myc. c-myc... |
| Clonality | Monoclonal |
| Clone Designation | [MYC/7150R] |
| Molecular Weight | 62-64kDa |
| NCBI | 4609 |

| | |
|-------------------|--|
| UniProt | P01106 |
| Form | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Recombinant Monoclonal Antibody |
| Application Notes | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes),Optimal dilutio |