

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

**Telephone:** +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## **Product Datasheet**

## FOXP3 (Forkhead Box Protein P3) / Scurfin Antibody, IgG1, Clone: [FXP3/197], Mouse, Monoclonal NBT-50943-MSM2-P0

| Article Name               | FOXP3 (Forkhead Box Protein P3) / Scurfin Antibody, IgG1, Clone: [FXP3/197], Mouse, Monoclonal   |
|----------------------------|--|
| Biozol Catalog Number      | NBT-50943-MSM2-P0  |
| Supplier Catalog Number    | 50943-MSM2-P0  |
| Alternative Catalog Number | NBT-50943-MSM2-P0-20,NBT-50943-MSM2-P0-100   |
| Manufacturer               | NeoBiotechnologies   |
| Host                       | Mouse  |
| Category                   | Antikörper   |
| Application                | FC, IHC, WB  |
| Species Reactivity         | Human, Monkey, Mouse   |
| Immunogen                  | Recombinant human full-length FOXP3 protein  |
| Product Description        | Recognizes a protein of 47-55kDa, which is identified as FOXP3. Its precise epitope is not known, but it has been mapped to the N-terminal portion of the protein. The FOX family of transcription factors is a large group of proteins that share a commo |
| Clonality                  | Monoclonal   |
| Clone Designation          | [FXP3/197]   |
| Molecular Weight           | 47-55kDa   |
| Isotype                    | IgG1   |

| NCBI              | 50943  |
|-------------------|--|
| UniProt           | Q9ZS1  |
| Form              | 200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double are a side. Also available WITHOUT BSA & Double at 1.0mg/ml.  |
| Antibody Type     | Monoclonal Antibody  |
| Application Notes | Flow Cytometry (0.5-1ug/million cells), Immunofluorescence (0.5-1ug/ml),,Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 |