

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

**Telephone:** +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## **Product Datasheet**

## Recombinant FOXP3 (Forkhead Box Protein P3) / Scurfin Antibody, Clone: [FOXP3/8015R], Rabbit, Monoclonal NBT-50943-RBM6-P0

| Article Name               | Recombinant FOXP3 (Forkhead Box Protein P3) / Scurfin Antibody, Clone: [FOXP3/8015R], Rabbit, Monoclonal   |
|----------------------------|--|
| Biozol Catalog Number      | NBT-50943-RBM6-P0  |
| Supplier Catalog Number    | 50943-RBM6-P0  |
| Alternative Catalog Number | NBT-50943-RBM6-P0-20,NBT-50943-RBM6-P0-100   |
| Manufacturer               | NeoBiotechnologies   |
| Host                       | Rabbit   |
| Category                   | Antikörper   |
| Application                | IHC  |
| Species Reactivity         | Human  |
| Immunogen                  | Recombinant fragment (around aa300-431) of human FOXP3 protein (exact sequence is proprietary)   |
| Product Description        | Recognizes a protein of 47-55kDa, which is identified as FOXP3. Its precise epitope is not known, but it has been mapped to the N-terminal portion of the protein. The FOX family of transcription factors is a large group of proteins that share a commo |
| Clonality                  | Monoclonal   |
| Clone Designation          | [FOXP3/8015R]  |
| Molecular Weight           | 47-55kDa   |
| NCBI                       | 50943  |

| UniProt           | Q9BZS1   |
|-------------------|--|
| Form              | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Protein |
| Antibody Type     | Recombinant Monoclonal Antibody  |
| Application Notes | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95&degC followed by cooling at RT for 20 minutes),Optimal dilutio   |