

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## **Product Datasheet**

## CD31 / PECAM-1 (Endothelial Cell Marker) Antibody, IgG1, Clone: [PECAM1/3529], Mouse, Monoclonal NBT-5175-MSM19-P1ABX

| Article Name               | CD31 / PECAM-1 (Endothelial Cell Marker) Antibody, IgG1, Clone:<br>[PECAM1/3529], Mouse, Monoclonal   |
|----------------------------|---|
| Biozol Catalog Number      | NBT-5175-MSM19-P1ABX  |
| Supplier Catalog Number    | 5175-MSM19-P1ABX  |
| Alternative Catalog Number | NBT-5175-MSM19-P1ABX-100  |
| Manufacturer               | NeoBiotechnologies  |
| Host                       | Mouse   |
| Category                   | Antikörper  |
| Application                | IHC, WB   |
| Species Reactivity         | Human   |
| Immunogen                  | Recombinant fragment (around aa 625-738) of human CD31 protein (exact sequence is proprietary)  |
| Product Description        | CD31 (PECAM-1) is a transmembrane glycoprotein member of the<br>immunoglobulin supergene family of adhesion molecules. CD31 is<br>expressed by stem cells of the hematopoietic system and is primarily<br>used to identify and concentrate these cells for experime |
| Clonality                  | Monoclonal  |
| Clone Designation          | [PECAM1/3529]   |
| Molecular Weight           | $\sim$ 100kDa (endothelium) and $\sim$ 130kDa (platelets)   |
| Isotype                    | lgG1  |

| NCBI              | 5175  |
|-------------------|---|
| UniProt           | P16284  |
| Form              | 200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G.<br>Prepared in 10mM PBS with 0.05% BSA & amp; 0.05% azide. Also<br>available WITHOUT BSA & amp; azide at 1.0mg/ml.  |
| Antibody Type     | Monoclonal Antibody   |
| Application Notes | Western Blot (1-2ug/ml),,Immunohistochemistry (Formalin-fixed) (1-<br>2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues<br>requires boiling tissue sections in 10mM citrate buffer, pH 6.0, for 10-<br>20 min followed by cooling at RT for 20 minutes),Op |