

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

BOB.1 (B-Cell Marker) Antibody, Clone: [BOB1/7468], Mouse, Monoclonal NBT-5450-MSM9-P0

| | |
|----------------------------|---|
| Article Name | BOB.1 (B-Cell Marker) Antibody, Clone: [BOB1/7468], Mouse, Monoclonal |
| Biozol Catalog Number | NBT-5450-MSM9-P0 |
| Supplier Catalog Number | 5450-MSM9-P0 |
| Alternative Catalog Number | NBT-5450-MSM9-P0-20,NBT-5450-MSM9-P0-100 |
| Manufacturer | NeoBiotechnologies |
| Host | Mouse |
| Category | Antikörper |
| Application | IHC |
| Species Reactivity | Human |
| Immunogen | Recombinant fragment (around aa 148-255) of human BOB1 (POU2AF1) protein (exact sequence is proprietary) |
| Product Description | BOB.1 expression in a variety of established B-cell lines, representing different stages of B-cell development, has suggested a constitutive, B-cell-specific expression pattern. LP cells in nodular lymphocyte predominant Hodgkin lymphoma, because the... |
| Clonality | Monoclonal |
| Clone Designation | [BOB1/7468] |
| Molecular Weight | 35kDa |
| NCBI | 5450 |

| | |
|-------------------|--|
| UniProt | Q16633 |
| Form | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Monoclonal Antibody |
| Application Notes | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 min at RT),(Staining of formalin-fixed tissues requires boiling tissue sections in 10mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes),Optimal dilution for a specific |