

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

**Telephone:** +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## **Product Datasheet**

## DOG-1 / ANO1 (Marker for Gastrointestinal Stromal Tumors) Antibody, IgG1, Clone: [DG1/447 + DOG-1.1], Mouse, Monoclonal NBT-55107-MSM3-P1ABX

| Article Name               | DOG-1 / ANO1 (Marker for Gastrointestinal Stromal Tumors) Antibody, IgG1, Clone: [DG1/447 + DOG-1.1], Mouse, Monoclonal  |
|----------------------------|--|
| Biozol Catalog Number      | NBT-55107-MSM3-P1ABX   |
| Supplier Catalog Number    | 55107-MSM3-P1ABX   |
| Alternative Catalog Number | NBT-55107-MSM3-P1ABX-100   |
| Manufacturer               | NeoBiotechnologies   |
| Host                       | Mouse  |
| Category                   | Antikörper   |
| Application                | IHC  |
| Species Reactivity         | Human  |
| Immunogen                  | Recombinant human DOG-1 protein (DG1/447) + A synthetic peptide from human DOG-1 protein (MSDFVDWVIPDIPKDISQQIHKEKVLMVELFMREEQDKQQLL-ETCMEKERQKDEPPCNHHNTKACPDSLGSP-APSHAYHGGVL), conjugated to a carrier protein (DOG-1.1).                               |
| Product Description        | Expression of DOG-1 protein is elevated in the gastrointestinal stromal tumors (GISTs), c-kit signaling-driven mesenchymal tumors of the GI tract. DOG-1 is rarely expressed in other soft tissue tumors, which, due to appearance, may be difficult to di |
| Clonality                  | Monoclonal   |
| Clone Designation          | [DG1/447 + DOG-1.1]  |

| Molecular Weight  | ~114kDa  |
|-------------------|--|
| Isotype           | lgG1   |
| NCBI              | 55107  |
| UniProt           | Q5XX6  |
| Form              | 200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Protein |
| Antibody Type     | Monoclonal Antibody  |
| Application Notes | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes), Optimal dilution f  |