

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Recombinant Prostate Specific Acid Phosphatase (PSAP) Antibody, Clone: [ACPP/4495R], Rabbit, Monoclonal NBT-55-RBM5-P1ABX

| | |
|----------------------------|---|
| Article Name | Recombinant Prostate Specific Acid Phosphatase (PSAP) Antibody, Clone: [ACPP/4495R], Rabbit, Monoclonal |
| Biozol Catalog Number | NBT-55-RBM5-P1ABX |
| Supplier Catalog Number | 55-RBM5-P1ABX |
| Alternative Catalog Number | NBT-55-RBM5-P1ABX-100 |
| Manufacturer | NeoBiotechnologies |
| Host | Rabbit |
| Category | Antikörper |
| Application | IHC |
| Species Reactivity | Human |
| Immunogen | Recombinant fragment (around aa1-100) of human ACPP (exact sequence is proprietary) |
| Product Description | Prostatic Acid Phosphatase (PSAP or PAP) is a member of the histidine acid phosphatase family. It is a non-specific phosphatase that is capable of dephosphorylating tyrosine residues as well as phospholipids under mildly acidic conditions. Prostatic ... |
| Clonality | Monoclonal |
| Clone Designation | [ACPP/4495R] |
| Molecular Weight | 45kDa |
| NCBI | 55 |

| | |
|-------------------|--|
| UniProt | P15309 |
| Form | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Recombinant Monoclonal Antibody |
| Application Notes | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes),Optimal dilutio |