

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Recombinant PMEPA1 / TMEPAI (Tumor Suppressor Oncoprotein) Antibody, IgG1, Clone: [rPMEPA1/6422], Mouse, Monoclonal NBT-56937-MSM12-P0

Article Name	Recombinant PMEPA1 / TMEPAI (Tumor Suppressor Oncoprotein) Antibody, IgG1, Clone: [rPMEPA1/6422], Mouse, Monoclonal
Biozol Catalog Number	NBT-56937-MSM12-P0
Supplier Catalog Number	56937-MSM12-P0
Alternative Catalog Number	NBT-56937-MSM12-P0-20,NBT-56937-MSM12-P0-100
Manufacturer	NeoBiotechnologies
Host	Mouse
Category	Antikörper
Application	IHC
Species Reactivity	Human
Immunogen	Recombinant full-length human PMEPA1 protein
Product Description	PMEPA1 (prostate transmembrane protein, androgen induced 1 is a 287 amino acid single-pass membrane protein that contains WW-binding motifs and localizes to the cell membrane. Expressed at high levels in prostate, kidney and ovary, PMEPA1 interacts w
Clonality	Monoclonal
Clone Designation	[rPMEPA1/6422]
Molecular Weight	32kDa
Isotype	IgG1

NCBI	56937
UniProt	Q96W9
Form	200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double are applied to the substitution of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double are applied to the substitution of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double are applied to the substitution of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double are applied to the substitution of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double are applied to the substitution of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double are applied to the substitution of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double are applied to the substitution of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double are applied to the substitution of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double are applied to the substitution of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double are applied to the substitution of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double are applied to the substitution of Ab purified from Bioreactor Concentrate by Protein A/G. P
Antibody Type	Recombinant Monoclonal Antibody
Application Notes	Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes),Optimal dilutio