

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

**Telephone:** +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## **Product Datasheet**

## CD45RO (T-Cell Marker) Antibody, IgG2a, Clone: [UCHL-1 + T200/797], Mouse, Monoclonal NBT-5788-MSM19-P0

Article Name	CD45RO (T-Cell Marker) Antibody, IgG2a, Clone: [UCHL-1 + T200/797], Mouse, Monoclonal
Biozol Catalog Number	NBT-5788-MSM19-P0
Supplier Catalog Number	5788-MSM19-P0
Alternative Catalog Number	NBT-5788-MSM19-P0-20,NBT-5788-MSM19-P0-100
Manufacturer	NeoBiotechnologies
Host	Mouse
Category	Antikörper
Application	IHC, WB
Species Reactivity	Human
Immunogen	Cultured human T-cells from an IL-2-dependent T-cell line (CA1) (UCHL-1), Human T-lymphocytes (T200/797)
Product Description	Recognizes a 180-185kDa protein, identified as isoform of leukocyte common antigen (CD45RO). This antibody reacts with mature activated T-cells, most thymocytes, and a sub-population of resting T-cells within both CD4 and CD8 subsets. It shows no rea
Clonality	Monoclonal
Clone Designation	[UCHL-1 + T200/797]
Molecular Weight	180-185kDa
Isotype	IgG2a

NCBI	5788
UniProt	P08575
Form	200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double are also available WITHOUT BSA & Double at 1.0mg/ml.
Antibody Type	Monoclonal Antibody
Application Notes	Western Blot (1-2ug/ml), Immunohistochemistry (Formalin-fixed) (0.25-0.5ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 2