

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

**Telephone:** +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## **Product Datasheet**

## CD45 / LCA (B-Cell Marker) Antibody, IgG1, Clone: [PTPRC/1666], Mouse, Monoclonal NBT-5788-MSM35-P1ABX

| Article Name               | CD45 / LCA (B-Cell Marker) Antibody, IgG1, Clone: [PTPRC/1666], Mouse, Monoclonal  |
|----------------------------|--|
| Biozol Catalog Number      | NBT-5788-MSM35-P1ABX   |
| Supplier Catalog Number    | 5788-MSM35-P1ABX   |
| Alternative Catalog Number | NBT-5788-MSM35-P1ABX-100   |
| Manufacturer               | NeoBiotechnologies   |
| Host                       | Mouse  |
| Category                   | Antikörper   |
| Application                | FC, WB   |
| Species Reactivity         | Human  |
| Immunogen                  | Human recombinant full-length CD45 protein   |
| Product Description        | CD45R, also designated CD45 and PTPRC, has been identified as a transmembrane glycoprotein, broadly expressed among hematopoietic cells. Multiple isoforms of CD45R are distributed throughout the immune system according to cell type. These isoforms ar |
| Clonality                  | Monoclonal   |
| Clone Designation          | [PTPRC/1666]   |
| Molecular Weight           | 180-220kDa   |
| Isotype                    | lgG1   |

| NCBI              | 5788   |
|-------------------|--|
| UniProt           | P08575   |
| Form              | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prote |
| Antibody Type     | Monoclonal Antibody  |
| Application Notes | Flow Cytometry (1-2ug/million cells), Western Blot (1-2ug/ml),,Optimal dilution for a specific application should be determined.   |