

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Bcl-2 (Apoptosis & Follicular Lymphoma Marker) Antibody, IgG1, Clone: [100/D5+124], Mouse, Monoclonal NBT-596-MSM4-P0

Article Name	Bcl-2 (Apoptosis & Follicular Lymphoma Marker) Antibody, IgG1, Clone: [100/D5+124], Mouse, Monoclonal
Biozol Catalog Number	NBT-596-MSM4-P0
Supplier Catalog Number	596-MSM4-P0
Alternative Catalog Number	NBT-596-MSM4-P0-20,NBT-596-MSM4-P0-100
Manufacturer	NeoBiotechnologies
Host	Mouse
Category	Antikörper
Application	FC, IF, IHC, WB
Species Reactivity	Human
Immunogen	A synthetic peptide, aa41-54 (GAAPAPGIFSSQPG-Cys) of human Bcl-2 (100/D5 & 124)
Product Description	This antibody recognizes a protein of 25-26kDa, identified as the bcl-2 lpha oncoprotein. It shows no cross-reaction with Bcl-x or Bax protein. Expression of bcl-2 lpha oncoprotein inhibits the programmed cell death (apoptosis). In most follicular ly
Clonality	Monoclonal
Clone Designation	[100/D5+124]
Molecular Weight	25-26kDa
Isotype	IgG1

NCBI	596
UniProt	P10415
Form	200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double are a side. Also available WITHOUT BSA & Double at 1.0mg/ml.
Antibody Type	Monoclonal Antibody
Application Notes	Flow Cytometry (1-2ug/million cells), Immunofluorescence (1-2ug/ml), Western Blot (1-2ug/ml), Immunohistochemistry (Formalinfixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues is enhanced by heating tissue sections in 10mM Tris wit