

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

BcI-X (Apoptosis Marker) Antibody, IgG2a, Clone: [SPM165], Mouse, Monoclonal NBT-598-MSM2X-P1ABX

Article Name	Bcl-X (Apoptosis Marker) Antibody, IgG2a, Clone: [SPM165], Mouse, Monoclonal
Biozol Catalog Number	NBT-598-MSM2X-P1ABX
Supplier Catalog Number	598-MSM2X-P1ABX
Alternative Catalog Number	NBT-598-MSM2X-P1ABX-100
Manufacturer	NeoBiotechnologies
Host	Mouse
Category	Antikörper
Application	FC, IF, IHC, WB
Species Reactivity	Human, Mouse, Porcine, Rat
Immunogen	A synthetic peptide, aa 3-14 (Cys-QSNRELVVDFLS) of human Bcl-X protein
Product Description	Recognizes a protein of 27kDa, identified as the Bcl-X protein. This MAb shows no cross-reaction with Bcl-2 or Bax protein. Bcl-X has two isoforms, Bcl-XL (long), a 241 amino acid protein which suppresses cell death. And Bcl-XS (short), a 178 amino a
Clonality	Monoclonal
Clone Designation	[SPM165]
Molecular Weight	27kDa
Isotype	IgG2a

NCBI	598
UniProt	Q07817
Form	200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double are a side. Also available WITHOUT BSA & Double at 1.0mg/ml.
Antibody Type	Monoclonal Antibody
Application Notes	Flow Cytometry (1-2ug/million cells), Immunofluorescence (1-2ug/ml), Western Blot (1-2ug/ml), Immunohistochemistry (Formalinfixed) (1-2ug/ml for 30 minutes at RT) (Staining of formalin-fixed tissues requires heating tissue sections in 1mM EDTA buffer, p