

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Recombinant Bcl-X (Apoptosis Marker) Antibody, IgG2a, Clone: [rBCL2L1/4508], Mouse, Monoclonal NBT-598-MSM8-P0

Article Name	Recombinant Bcl-X (Apoptosis Marker) Antibody, IgG2a, Clone: [rBCL2L1/4508], Mouse, Monoclonal
Biozol Catalog Number	NBT-598-MSM8-P0
Supplier Catalog Number	598-MSM8-P0
Alternative Catalog Number	NBT-598-MSM8-P0-20,NBT-598-MSM8-P0-100
Manufacturer	NeoBiotechnologies
Host	Mouse
Category	Antikörper
Application	IHC
Species Reactivity	Human
Immunogen	Recombinant fragment (around aa1-100) of human BCL2L1 protein (exact sequence is proprietary)
Product Description	Recognizes a protein of 27kDa, identified as the Bcl-X protein. This MAb shows no cross-reaction with Bcl-2 or Bax protein. Bcl-X has two isoforms, Bcl-XL (long), a 241-amino acid protein which suppresses cell death. And Bcl-XS (short), a 178-amino a
Clonality	Monoclonal
Clone Designation	[rBCL2L1/4508]
Molecular Weight	27kDa
Isotype	IgG2a

NCBI	598
UniProt	Q07817
Form	200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double are a side. Also available WITHOUT BSA & Double at 1.0mg/ml.
Antibody Type	Recombinant Monoclonal Antibody
Application Notes	Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95 °C followed by cooling at RT for 20 minutes),Optimal diluti