

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **Recombinant CD269 / TNFRSF17 / BCMA (B-Cell Maturation Protein) Antibody, Clone: [CD269/8190R], Rabbit, Monoclonal NBT-608-RBM8-P0**

|                            |   |
|----------------------------|---|
| Article Name               | Recombinant CD269 / TNFRSF17 / BCMA (B-Cell Maturation Protein) Antibody, Clone: [CD269/8190R], Rabbit, Monoclonal  |
| Biozol Catalog Number      | NBT-608-RBM8-P0   |
| Supplier Catalog Number    | 608-RBM8-P0   |
| Alternative Catalog Number | NBT-608-RBM8-P0-20,NBT-608-RBM8-P0-100  |
| Manufacturer               | NeoBiotechnologies  |
| Host                       | Rabbit  |
| Category                   | Antikörper  |
| Application                | ELISA, IHC  |
| Species Reactivity         | Human   |
| Immunogen                  | Recombinant human CD269 protein fragment (around aa 1-184) (exact sequence is proprietary)  |
| Product Description        | The B cell maturation protein (BCMA) is a type I integral membrane protein that belongs to the tumor necrosis factor receptor (TNF-R) superfamily. It is expressed as a 184 amino acid peptide that is expressed only in mature B-lymphocytes and is locat... |
| Clonality                  | Monoclonal  |
| Clone Designation          | [CD269/8190R]   |
| Molecular Weight           | 20kDa   |
| NCBI                       | <a href="#">608</a>   |

|                   |  |
|-------------------|--|
| UniProt           | <a href="#">Q02223</a>   |
| Form              | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.   |
| Antibody Type     | Recombinant Monoclonal Antibody  |
| Application Notes | ELISA (For coating, order antibody without BSA), Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95&degC followed |