

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

**Telephone:** +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## **Product Datasheet**

## Recombinant CD71 / Transferrin Receptor (TFRC) (Extracellular Domain) Antibody, Clone: [TFRC/2898R], Rabbit, Monoclonal NBT-7037-RBM10-P0

Article Name	Recombinant CD71 / Transferrin Receptor (TFRC) (Extracellular Domain) Antibody, Clone: [TFRC/2898R], Rabbit, Monoclonal
Biozol Catalog Number	NBT-7037-RBM10-P0
Supplier Catalog Number	7037-RBM10-P0
Alternative Catalog Number	NBT-7037-RBM10-P0-20,NBT-7037-RBM10-P0-100
Manufacturer	NeoBiotechnologies
Host	Rabbit
Category	Antikörper
Application	IHC
Species Reactivity	Human
Immunogen	Recombinant human extracellular TFRC protein fragment (aa94-212) (exact sequence is proprietary)
Product Description	It recognizes a $\sim 90\text{-}95\text{kDa}$ protein which is identified as cell surface transferrin receptor (CD71), a disulfide-bonded homodimeric glycoprotein of 180-190kDa. This MAb is highly specific to CD71 and shows no cross-reaction with other related proteins
Clonality	Monoclonal
Clone Designation	[TFRC/2898R]
Molecular Weight	85-95kDa (monomer), 190kDa (dimer)
NCBI	7037

UniProt	P02786
Form	200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double Concentrate by Protein A/G. Prote
Antibody Type	Recombinant Monoclonal Antibody
Application Notes	Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes),Optimal dilution fo