

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Recombinant Topoisomerase II alpha (Proliferation & Drug-Resistance Marker) Antibody, Clone: [TOP2A/4397R], Rabbit, Monoclonal NBT-7153-RBM3-P1ABX

| | |
|----------------------------|---|
| Article Name | Recombinant Topoisomerase II alpha (Proliferation & Drug-Resistance Marker) Antibody, Clone: [TOP2A/4397R], Rabbit, Monoclonal |
| Biozol Catalog Number | NBT-7153-RBM3-P1ABX |
| Supplier Catalog Number | 7153-RBM3-P1ABX |
| Alternative Catalog Number | NBT-7153-RBM3-P1ABX-100 |
| Manufacturer | NeoBiotechnologies |
| Host | Rabbit |
| Category | Antikörper |
| Application | IHC |
| Species Reactivity | Human |
| Immunogen | Synthetic peptide corresponding to Topo II alpha residues within aa1431-1531 of Topo II alpha |
| Product Description | The topoisomerase II- isoform is a 170kDa nuclear protein and plays an important role in DNA synthesis and RNA transcription, as well as chromosomal segregation during mitosis. Topoisomerase II- is reported to be a sensitive and specific marker of la... |
| Clonality | Monoclonal |
| Clone Designation | [TOP2A/4397R] |
| Molecular Weight | 170kDa |

| | |
|-------------------|--|
| NCBI | 7153 |
| UniProt | P11388 |
| Form | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Recombinant Monoclonal Antibody |
| Application Notes | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes),Optimal dilutio |