

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Tryptase (Mast Cell Marker) Antibody, IgG1, Clone: [TPSAB1/1963], Mouse, Monoclonal NBT-7177-MSM3-P0

| Article Name | Tryptase (Mast Cell Marker) Antibody, IgG1, Clone: [TPSAB1/1963], Mouse, Monoclonal |
|----------------------------|--|
| Biozol Catalog Number | NBT-7177-MSM3-P0 |
| Supplier Catalog Number | 7177-MSM3-P0 |
| Alternative Catalog Number | NBT-7177-MSM3-P0-20,NBT-7177-MSM3-P0-100 |
| Manufacturer | NeoBiotechnologies |
| Host | Mouse |
| Category | Antikörper |
| Application | IHC |
| Species Reactivity | Human |
| Immunogen | Recombinant human Tryptase protein fragment (around aa 115-233) (Exact sequence is proprietary) |
| Product Description | Tryptases comprise a family of trypsin-like serine proteases (peptidase family S1). Tryptases are stored in mast cell secretory granules and basophils. Mast cells are connective tissue cells derived from blood-forming tissues that line arterial walls |
| Clonality | Monoclonal |
| Clone Designation | [TPSAB1/1963] |
| Molecular Weight | 31-36kDa |
| Isotype | lgG1 |

| NCBI | 7177 |
|-------------------|--|
| UniProt | Q15661 |
| Form | 200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & Double are a side. Also available WITHOUT BSA & Double at 1.0mg/ml. |
| Antibody Type | Monoclonal Antibody |
| Application Notes | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes), Optimal dilution |