

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Carbonic Anhydrase VIII Antibody, IgG2a, Clone: [CA8/6572], Mouse, Monoclonal NBT-767-MSM2-P0

| | |
|----------------------------|---|
| Article Name | Carbonic Anhydrase VIII Antibody, IgG2a, Clone: [CA8/6572], Mouse, Monoclonal |
| Biozol Catalog Number | NBT-767-MSM2-P0 |
| Supplier Catalog Number | 767-MSM2-P0 |
| Alternative Catalog Number | NBT-767-MSM2-P0-20,NBT-767-MSM2-P0-100 |
| Manufacturer | NeoBiotechnologies |
| Host | Mouse |
| Category | Antikörper |
| Application | IHC |
| Species Reactivity | Human |
| Immunogen | Recombinant full-length human CA8 protein |
| Product Description | The protein encoded by this gene was initially named CA-related protein because of sequence similarity to other known carbonic anhydrase genes. However, the gene product lacks carbonic anhydrase activity (i.e., the reversible hydration of carbon diox... |
| Clonality | Monoclonal |
| Clone Designation | [CA8/6572] |
| Molecular Weight | 32.97kDa |
| Isotype | IgG2a |

| | |
|-------------------|---|
| NCBI | 767 |
| UniProt | P35219 |
| Form | 200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml. |
| Antibody Type | Monoclonal Antibody |
| Application Notes | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95 °C followed by cooling at RT for 20 minutes),Optimal diluti |